arXiv:1704.04106v1 [astro-ph.EP] 13 Apr 2017

Molecular line shape parameters for exoplanetary atmospheric applications

Sergey Yurchenko, Jonathan Tennyson and Emma J. Barton

Department of Physics and Astronomy, University College London, Gower Street, WC1E 6BT London, UK

E-mail: s.yurchenko@ucl.ac.uk

Abstract.

We describe the recent updates to the ExoMol database regarding the molecular spectral line shapes. ExoMol provides comprehensive molecular line lists with a special emphasis on the applications involving characterization of atmospheres of exoplanets and cool stars. Among important requirements of such applications are (i) the broadening parameters for hydrogen and helium dominating atmospheres and (ii) very broad ranges of temperature and pressures. The current status of the available line shape data in the literature, demands from the exoplanetary community and their specific needs are discussed.

1. Introduction

Molecules in the atmospheres of exoplanets, and similar astronomical objects such as brown dwarfs, are observed at pressures approaching atmospheric and therefore display pronounced pressure effects in their spectra [1, 2]. There is, however, a severe lack of data appropriate for reproducing these pressure effects.

The ExoMol project provide extensive line lists for a large range of molecules, mostly related to the atmospheric retrievals for exoplanets and cool stars [3]. The selection of molecules and the spectroscopic coverage in ExoMol are dictated by the requirements from these applications. A major requirement is that the line lists must be sufficiently complete to be applicable for high temperatures specific for atmospheres of most of these objects [4]. Spectroscopic data for almost 40 molecules are currently covered by the ExoMol database, including diatomic, triatomic, tetratomic and two pentatomic species. The available line lists were generated both by ExoMol and other groups, see Table 1.

Recently we have undertaken a major upgrade of the ExoMol database [5]. The upgrade was motivated by the growing role of ExoMol as a major provider of the hot spectroscopic line lists for atmospheric retrievals from exoplanetary observations. The most important modifications include: (i) introduction of application programming interface (API); (ii) line shape parameters; (iii) cooling functions; (iv) radiative life times [6] and Landé g-factors [7]. In this contribution the new line shape data provided by ExoMol are discussed.

2. Requirements for the line shape parameters from the exoplanetary applications.

It is well recognized that Voigt profiles only provide an approximate solution, especially for high resolution atmospheric studies [8]. However Voigt profiles are in widespread use and are

easily computed [9]; they are therefore uniformly used to represent pressure broadening effects in exoplanetary models and hence in the ExoMol database.

Table 1 illustrates the coverage of the broadening parameters in ExoMol, which is also a reflection of the current status in the field. Below we list the main factors affecting the development of the line shapes for the exoplanetary applications.

(i) The dominant species in gas giant planets such as hot Jupiter atmospheres are H_2 and He, often at high pressure [10]. However most of the line shapes parameters available in the literature are for the air (or N_2) as the main broadener and motivated by the terrestrial applications. Even though the needs for the line shapes broadened by H_2 and He have been recently recognized by HITRAN [11], the corresponding line shape data is still incomplete [12, 2].

(ii) The large size of the line lists in ExoMol, which consist of tens of billions of lines for larger molecules, make it practically impossible to populate the whole database with accurate, line-by-line line shape parameters, experimental or theoretical. Only simple and very approximate models (e.g. based only on one quantum number J'') can afford productions of the data on such a large scale. However even simple models do not exist for most of the species in question, especially for H_2 and He as broadeners.

(iii) The high temperatures ($T > 1000K$) of hot Jupiters, and other exoplanets of current interest [13], require line profiles for a large range of rotation and vibration excitations over extended temperature ranges. It is considered to be important to model at least the rotational (J) dependence of the line widths broadened by H_2 [10].

(iv) There is no agreement on the value of the line wing cut-off to be used when computing Voigt profiles. On one hand, it is imperative to use a reasonable cut-off distance in order to speed up the line-by-line calculations of the molecular opacities. On the other hand, the choice of the cut-off is known to effect the results at high pressures. Common practices include a fixed cut-off (e.g. 25 cm^{-1} or 100 cm^{-1} [10]), variable cut-off depending on pressure P (e.g. min(25P; 100) cm^{-1} [14]) or proportional to the line (half-)width (e.g. 500 widths [15]). The importance of folding back the truncated wings to ensure that the strength of the profile is conserved [14] is also considered in certain circumstances.

(v) Efficient computational algorithms for the Voigt evaluations [9, 16] as well as the sampling method [17] are required when billions of lines are involved.

In practice it is computationally prohibitive to perform radiative transport calculations on hot exoplanets line-by-line. Therefore the models either use precomputed cross sections, such as is done by τ-Rex [37, 123], or tables of k-coefficients, as done by the NEMISIS [124]. This means that appropriate values are precomputed on a grid of temperatures and pressures as inputs to such codes.

3. Conclusion

We are in the advanced stages of developing a diet for pressure broadening of molecules present in exoplanetary atmospheres and those of other hot astronomical objects. A full discussion of this problem will be given elsewhere [125].

Pressure broadening data is urgently needed for atmospheric studies (retrievals) of exoplanets and cool stars. There is huge demand on the comprehensive solutions of the line shape problems for most of the molecules important for exoplanetary studies. The ExoMol database is arguably the main source of opacities for hot species important for modeling these atmospheres, where we have created a structure for depositing and curating any molecular data important for spectroscopic properties of hot atmospheric and other gaseous environments. We invite the molecular data producers to contribute to this database. The line shape data is especially

Table 1. Molecular line lists included in the ExoMol database and Voigt line shape parameters when available.

Molecule	Reference	Broadener	Molecule	Reference	Broadener
AlO	[18]		HD^+	[19]	
BeH	[20]		HeH^+	[21]	
CaH	[20]		HNO_3	[22]	Air
CaH	[23]		KCl	[24]	
CaO	[25]		LiH	[19]	
CH	[26]		LiH^+	[19]	
CH_4	[27]	Air, He, H_2, CH_4	MgH	[20]	
CN	[28]		MgH	[29]	
CP	[30]		NaCl	[24]	
CrH	[31]		NaH	[32]	
CS	[33]	Air, CS	NH	[34]	
FeH	[35]		NH_3	[36]	Air, H_2, He, NH_3
H_2CO	[37]	Air, He, H_2, H_2CO	PH_3	[37]	Air, He, H_2, PH_3
H_2O	[38]	Air, H_2, He, H_2O	PN	[39]	
H_3^+	[40]		ScH	[41]	
HCl	[42]	Air, H_2, He, HCl	SiO	[43]	
HCN/HNC	[44]	Air, He, H_2, HCN	TiH	[45]	
SO_2	[46]	Air, He, H_2	SO_3	[47]	
H_2S	[48]		H_2O_2	[49]	
VO	[50]				

important as the line parameters are missing or incomplete for the H_2-rich atmospheres even for the most important absorbers. Exoplanetary atmospheric retrieval is a hot topic at the moment with a lot of interest from the society, which makes it a good place to be for an expert in the molecular line profiles. The lack of data, a strong demand from the field and interest from the public is a very attractive mixture for work in this direction to be properly recognized and rewarded.

We also invite the community to visit and test the new ExoMol database at www.exomol.com. Any feedback will be greatly appreciated.

Acknowledgements
This work is supported by ERC Advanced Investigator Project 267219. We thank the support of the COST action MOLIM (CM1405).

References

[1] Tinetti G, Tennyson J, Griffiths C A and Waldmann I 2012 *Phil. Trans. Royal Soc. London A* **370** 2749–2764
[2] Freedman R S, Marley M S and Lodders K 2008 *Astrophys. J. Suppl.* **174** 504
[3] Tennyson J and Yurchenko S N 2012 *Mon. Not. R. Astron. Soc.* **425** 21–33
[4] Yurchenko S N, Tennyson J, Bailey J, Hollis M D J and Tinetti G 2014 *Proc. Nat. Acad. Sci.* **111** 9379–9383
[5] Tennyson J, Yurchenko S N, Al-Refaie A F, Barton E J, Chubb K L, Coles P A, Diamantopoulou S, Gorman M N, Hill C, Lam A Z, Lodi L, McKemmish L K, Na Y, Owens A, Polyansky O L, Rivlin T, Sousa-Silva C, Underwood D S, Yachmenev A and Zak E 2016 *J. Mol. Spectrosc.*
[6] Tennyson J, Hulme K, Naim O K and Yurchenko S N 2016 *J. Phys. B: At. Mol. Opt. Phys.* **49** 044002
[7] Semenov M, Yurchenko S N and Tennyson J 2016 *J. Mol. Spectrosc.*
[8] Tennyson J, Bernath P F, Campargue A, Császár A G, Daumont L, Gamache R R, Hodges J T, Lisak D, Naumenko O V, Rothman L S, Tran H, Zobov N F, Buldyreva J, Boone C D, De Vizia M D, Gianfrani

Table 2. Voigt parameters available in the literature and used in the ExoMol database.

Molecule	Broadener	Reference
H_2O	H_2	[51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63]
H_2O	He	[64, 65, 66, 67, 68, 51, 52, 53, 56, 57, 69, 60, 61, 62]
H_2O	Air	[70, 71, 72, 73, 74, 75]
H_2O	H_2O	[70, 71, 76, 77, 78]
CH_4	H_2	[79, 80, 81, 82]
CH_4	He	[83, 84, 85, 81]
CH_4	Air	[86, 87, 88]
CH_4	CH_4	[89, 90, 87]
NH_3	H_2	[91, 92, 14, 93]
NH_3	He	[91, 92, 14, 93]
NH_3	Air	[94, 95, 93]
NH_3	NH_3	[94, 96, 97, 98, 99, 100]
PH_3	H_2	[101, 96, 97, 99]
PH_3	He	[102, 96, 97, 99]
PH_3	Air/N_2	[102, 103, 104]
PH_3	PH_3	[103, 104, 105, 106]
H_2CO	H_2	[107]
H_2CO	He	[107]
H_2CO	N_2	[108]
H_2CO	H_2CO	[108]
SO_2	H_2	[109]
SO_2	He	[109, 110]
SO_2	Air	[109, 110]
HCN	Air/N_2	[111, 112, 113]
HCN	HCN	[112, 114]
HCN	H_2	[115]
HCN	He	[115]
HCl	N_2	[116]
HCl	HCl	[117]
HCl	H_2	[118]
HCl	He	[119, 116]
CS	Air	[120]
CS	CS	[121]
HNO_3	Air	[122]

L, Hartmann J M, McPheat R, Murray J, Ngo N H, Polyansky O L and Weidmann D 2014 *Pure Appl. Chem.* **86** 1931–1943

[9] Humlicek J 1979 *J. Quant. Spectrosc. Radiat. Transf.* **21** 309–313

[10] Amundsen D S, Baraffe I, Tremblin P, Manners J, Wolfgang H, Mayne N J and Acreman D M 2014 *Astron. Astrophys.* **564** A59

[11] Wilzewski J S, Gordon I E, Kochanov R V, Hill C and Rothman L S 2016 *J. Quant. Spectrosc. Radiat. Transf.* **168** 193 – 206

[12] Madhusudhan N 2012 *Astrophys. J.* **758** 36

[13] Tsiaras A, Rocchetto M, Waldmann I P, Tinetti G, Varley R, Morello G, Barton E J, Yurchenko S N and Tennyson J 2016 *Astrophys. J.* **820** 99

[14] Sharp C M and Burrows A 2007 *Astrophys. J. Suppl.* **168** 140

[15] Hedges C and Madhusudhan N 2016 *Mon. Not. R. Astron. Soc.* **458** 1427–1449

[16] Schreier F 2011 *J. Quant. Spectrosc. Radiat. Transf.* **112** 1010 – 1025 ISSN 0022-4073

[17] Evans T M, Sing D K, Wakeford H R, Nikolov N, Ballester G E, Drummond B, Kataria T, Gibson N P, Amundsen D S and Spake J 2016 *Astrophys. J.* **822** L4
[18] Patrascu A T, Tennyson J and Yurchenko S N 2015 *Mon. Not. R. Astron. Soc.* **449** 3613–3619
[19] Coppola C M, Lodi L and Tennyson J 2011 *Mon. Not. R. Astron. Soc.* **415** 487–493
[20] Yadin B, Vaness T, Conti P, Hill C, Yurchenko S N and Tennyson J 2012 *Mon. Not. R. Astron. Soc.* **425** 34–43
[21] Engel E A, Doss N, Harris G J and Tennyson J 2005 *Mon. Not. R. Astron. Soc.* **357** 471–477
[22] Pavlyuchko A I, Yurchenko S N and Tennyson J 2015 *Mon. Not. R. Astron. Soc.* **452** 1702–1706
[23] Li G, Harrison J J, Ram R S, Western C M and Bernath P F 2012 *J. Quant. Spectrosc. Radiat. Transf.* **113** 67 – 74 ISSN 0022-4073 URL `http://www.sciencedirect.com/science/article/pii/S0022407311003323`
[24] Barton E J, Chiu C, Golpayegani S, Yurchenko S N, Tennyson J, Frohman D J and Bernath P F 2014 *Mon. Not. R. Astron. Soc.* **442** 1821–1829
[25] Yurchenko S N, Blissett A, Asari U, Vasilios M, Hill C and Tennyson J 2016 *Mon. Not. R. Astron. Soc.* **456** 4524–4532
[26] Masseron T, Plez B, Van Eck S, Colin R, Daoutidis I, Godefroid M, Coheur P F, Bernath P, Jorissen A and Christlieb N 2014 *Astron. Astrophys.* **571** A47
[27] Yurchenko S N and Tennyson J 2014 *Mon. Not. R. Astron. Soc.* **440** 1649–1661
[28] Brooke J S A, Ram R S, Western C M, Li G, Schwenke D W and Bernath P F 2014 *Astrophys. J. Suppl.* **210** 23
[29] GharibNezhad E, Shayesteh A and Bernath P F 2013 *Mon. Not. R. Astron. Soc.* **432** 2043–2047
[30] Ram R S, Brooke J S A, Western C M and Bernath P F 2014 *J. Quant. Spectrosc. Radiat. Transf.* **138** 107–115
[31] Burrows A, Ram R S, Bernath P, Sharp C M and Milsom J A 2002 *Astrophys. J.* **577** 986–992
[32] Rivlin T, Lodi L, Yurchenko S N, Tennyson J and Le Roy R J 2015 *Mon. Not. R. Astron. Soc.* **451** 5153–5157
[33] Paulose G, Barton E J, Yurchenko S N and Tennyson J 2015 *Mon. Not. R. Astron. Soc.* **454** 1931–1939
[34] Brooke J S A, Ram R S, Western C M, Li G, Schwenke D W and Bernath P F 2014 *Astrophys. J. Suppl.* **210** 23 URL `http://stacks.iop.org/0067-0049/210/i=2/a=23`
[35] Wende S, Reiners A, Seifahrt A and Bernath P F 2010 *Astron. Astrophys.* **523** A58
[36] Yurchenko S N, Barber R J and Tennyson J 2011 *Mon. Not. R. Astron. Soc.* **413** 1828–1834
[37] Waldmann I P, Tinetti G, Barton E J, Yurchenko S N and Tennyson J 2015 *Astrophys. J.* **802** 107
[38] Barber R J, Tennyson J, Harris G J and Tolchenov R N 2006 *Mon. Not. R. Astron. Soc.* **368** 1087–1094
[39] Yorke L, Yurchenko S N, Lodi L and Tennyson J 2014 *Mon. Not. R. Astron. Soc.* **445** 1383–1391
[40] Neale L, Miller S and Tennyson J 1996 *Astrophys. J.* **464** 516–520
[41] Lodi L, Yurchenko S N and Tennyson J 2015 *Mol. Phys.* **113** 1559–1575
[42] Li G, Gordon I E, Bernath P F and Rothman L S 2011 *J. Quant. Spectrosc. Radiat. Transf.* **112** 1543 – 1550
[43] Barton E J, Yurchenko S N and Tennyson J 2013 *Mon. Not. R. Astron. Soc.* **434** 1469–1475
[44] Barber R J, Strange J K, Hill C, Polyansky O L, Mellau G C, Yurchenko S N and Tennyson J 2014 *Mon. Not. R. Astron. Soc.* **437** 1828–1835
[45] Burrows A, Dulick M, Bauschlicher C W, Bernath P F, Ram R S, Sharp C M and Milsom J A 2005 *Astrophys. J.* **624** 988–1002
[46] Underwood D S, Tennyson J, Yurchenko S N, Huang X, Schwenke D W, Lee T J, Clausen S and Fateev A 2016 *Mon. Not. R. Astron. Soc.* **459** 3890–3899
[47] Underwood D S, Tennyson J, Yurchenko S N, Clausen S and Fateev A 2016 *Mon. Not. R. Astron. Soc.* (submitted)
[48] Azzam A A A, Yurchenko S N, Tennyson J and Naumenko O V 2016 *Mon. Not. R. Astron. Soc.* **460** 4063–4074
[49] Al-Refaie A F, Polyansky O L, I R, Ovsyannikov, Tennyson J and Yurchenko S N 2016 *Mon. Not. R. Astron. Soc.* **461** 1012–1022
[50] McKemmish L K, Yurchenko S N and Tennyson J 2016 *Mon. Not. R. Astron. Soc.* submitted
[51] Lavrentieva N N, Voronin B A, Naumenko O V, Bykov A D and Fedorova A A 2014 *Icarus* **236** 38
[52] Barton E J, Hill C, Yurchenko S N, Tennyson J, Dudaryonok A and Lavrentieva N N 2016 *J. Quant. Spectrosc. Radiat. Transf.* (in preparation)
[53] Steyert D W, Wang W, Sirota J, Donahue N M and Reuter D C 2004 *J. Quant. Spectrosc. Radiat. Transf.* **83** 183–191
[54] Brown L R and Plymate C 1996 *J. Quant. Spectrosc. Radiat. Transf.* **56** 263 – 282
[55] Brown L R, Benner D C, Devi V M, Smith M A H and Toth R A 2005 *J. Mol. Struct.* **742** 111–122
[56] Gamache R R, Lynch R and Brown L R 1996 *J. Quant. Spectrosc. Radiat. Transf.* **56** 471 – 487
[57] Dick M J, Drouin B J and Pearson J C 2009 *J. Quant. Spectrosc. Radiat. Transf.* **110** 619 – 627

[58] Faure A, Wiesenfeld L, Tennyson J and Drouin B J 2013 *J. Quant. Spectrosc. Radiat. Transf.* **116** 79–86
[59] Langlois S, Birbeck T P and Hanson R K 1994 *J. Mol. Spectrosc.* **167** 272 – 281
[60] Dutta J M, Jones C R, Goyette T M and Lucia F C 1993 *Icarus* **102** 232–239
[61] Golubiatnikov Y G 2005 *J. Mol. Spectrosc.* **230** 196–198
[62] Zeninari V, Parvitte B, Courtois D, Lavrentieva N N, Ponomarev Y N and Durry G 2004 *Mol. Phys.* **102** 1697–1706
[63] Drouin B and Wiesenfeld L 2012 *Phys. Rev. A* **86** 022705
[64] Petrova T M, Solodov A M, Solodov A A and Starikov V I 2016 *J. Mol. Spectrosc.* **321** 50–58
[65] Petrova T M, Solodov A M, Solodov A A and Starikov V I 2013 *J. Quant. Spectrosc. Radiat. Transf.* **129** 241–253
[66] Petrova T M, Solodov A M, Solodov A A and Starikov V I 2012 *Mol. Phys.* **110** 1493–1503
[67] Solodov A M and Starikov V I 2008 *Opt. Spectrosc.* **105** 14–20
[68] Solodov A M and Starikov V I 2009 *Mol. Phys.* **104** 43–51
[69] Lazarev V V, Ponomarev Y N, Sumpf B, Fleischmann O, Waschull J, Kronfeldt H and Stroinova V N 1995 *J. Mol. Spectrosc.* **173** 177–193
[70] Mérienne M F, Jenouvrier A, Hermans C, Vandaele A C, Carleer M, Coheur P F, Colin R, Fally S and Bach M 2003 *J. Quant. Spectrosc. Radiat. Transf.* **82** 99–117
[71] Gamache R R and Hartmann J M 2004 *Can. J. Chem.* **82** 1013–1027
[72] Gasster S D, Townes C H, Goorvitch D and Valero F P J 1988 *J. Opt. Soc. Am. B* **5** 593–601
[73] Payne V, Delamere J, Cady-Pereira K, Gamache R, Moncet J L, Mlawer E and Clough S 2008 *IEEE Transactions on Geoscience and Remote Sensing* **46** 3601–3617
[74] Gamache R 2005 *J. Mol. Spectrosc.* **229** 9–18
[75] Gamache R and Laraia A 2009 *J. Mol. Spectrosc.* **257** 116–127
[76] Markov V 1994 *J. Mol. Spectrosc.* **164** 233–238
[77] Golubiatnikov G, Koshelev M and Krupnov A 2008 *J. Quant. Spectrosc. Radiat. Transf.* **109** 1828–1833
[78] Cazzoli G, Puzzarini C, Buffa G and Tarrini O 2008 *J. Quant. Spectrosc. Radiat. Transf.* **109** 2820–2831
[79] Pine A S 1992 *J. Chem. Phys.* **97** 773
[80] Margolis J S 1993 *J. Quant. Spectrosc. Radiat. Transf.* **50** 431
[81] Fox K, Jennings D E, Stern E A and Hunnard R 1988 *J. Quant. Spectrosc. Radiat. Transf.* **39** 473 – 476
[82] Strong K, Taylor F W, Calcutt S B, Remedios J J and Ballard J 1993 *J. Quant. Spectrosc. Radiat. Transf.* **50** 363 – 429
[83] Varanasi P and Chudamani S 1990 *J. Quant. Spectrosc. Radiat. Transf.* **43** 1
[84] Gabard T, Grigoriev I M, Grigorovich N M and Tonkov M V 2004 *J. Mol. Spectrosc.* **225** 123 – 131
[85] Grigoriev I M, Filippov N N, Tonkov M V, Gabard T and Le Doucen R 2001 *J. Quant. Spectrosc. Radiat. Transf.* **69** 189 – 204
[86] Smith M A H, Benner D C, Predoi-Cross A and Malathy Devi V 2009 *J. Quant. Spectrosc. Radiat. Transf.* **110** 639–653
[87] Brown L R, Benner D C, Champion J P, Devi V M, Fejard L, Gamache R R and et al 2003 *J. Quant. Spectrosc. Radiat. Transf.* **82** 219–238
[88] Antony B K, Niles D L, Wroblewski S B, Humphrey C M, Gabard T and Gamache R R 2008 *J. Mol. Spectrosc.* **251** 268–281
[89] Predoi-Cross A, Brown L R, Devi V M, Brawley-Tremblay M and Benner D C 2005 *J. Mol. Spectrosc.* **232** 231–246
[90] Smith M A H, Benner D C, Predoi-Cross A and Malathy Devi V 2009 *J. Quant. Spectrosc. Radiat. Transf.* **111** 1152–1166
[91] Pine A S, Markov V N, Buffa G and Tsrrini O 1993 *J. Quant. Spectrosc. Radiat. Transf.* **50** 337
[92] Hadded S, Arouli H, Orphal J, Bouanich J P and Hartmann J M 2001 *J. Mol. Spectrosc.* **210** 275
[93] Baldacchini G, D'Amato F, Buffa G, Tarrini O, Rosa M and Pelagalli F 2001 *Journal of Quantitative Spectroscopy and Radiative Transfer* **68** 625 – 633 ISSN 0022-4073 URL `http://www.sciencedirect.com/science/article/pii/S0022407300000431`
[94] Brown L R and Peterson D B 1994 *J. Mol. Spectrosc.* **168** 593–606
[95] Nemtchinov V, Sung K and Varanasi P 2004 *Journal of Quantitative Spectroscopy and Radiative Transfer* **83** 243 – 265 ISSN 0022-4073 URL `http://www.sciencedirect.com/science/article/pii/S0022407302003540`
[96] Levy A, Lacome N and Tarrago G 1993 *J. Mol. Spectrosc.* **157** 172 – 181
[97] Sergent-Rozey M, van Thanh N, Rossi I, Lacome N and Levy A 1988 *J. Mol. Spectrosc.* **131** 66 – 76
[98] Salem J, Bouanich J P, Walrand J, Aroui H and Blanquet G 2004 *J. Mol. Spectrosc.* **228** 23 – 30
[99] Pickett H M, Poynter R L and Cohen E A 1981 *J. Quant. Spectrosc. Radiat. Transf.* **26** 197 – 198
[100] Levy A, Lacome N and Tarrago G 1994 *J. Mol. Spectrosc.* **166** 20 – 31

[101] Bouanich J P, Salem J, Aroui H, Walrand J and Blanquet G 2004 *J. Quant. Spectrosc. Radiat. Transf.* **84** 195 – 205
[102] Salem J, Bouanich J P, Walrand J, Aroui H and Blanquet G 2005 *J. Mol. Spectrosc.* **232** 247 – 254
[103] Butler R A H, Sagui L, Kleiner I and Brown L R 2006 *J. Mol. Spectrosc.* **238** 178–192
[104] Kleiner I, Tarrago G, Cottaz C, Sagui L, Brown L R, Poynter R L, Pickett H M, Chen P, Pearson J C, Sams R L, Blake G A, Matsuura S, Nemtchinov V, Varanasi P, Fusina L and Di Lonardo G 2003 *J. Quant. Spectrosc. Radiat. Transf.* **82** 293 – 312
[105] Nguyen-Van-Thanh, Rossi I and Sergent-Rozey M 1989 *Journal of Molecular Spectroscopy* **135** 410 – 414 ISSN 0022-2852 URL http://www.sciencedirect.com/science/article/pii/0022285289901653
[106] Salem J, Aroui H, Bouanich J P, Walrand J and Blanquet G 2004 *Journal of Molecular Spectroscopy* **225** 174 – 181 ISSN 0022-2852 URL http://www.sciencedirect.com/science/article/pii/S002228520400075X
[107] Nerf R B J 1975 *J. Mol. Spectrosc.* **58** 451–473
[108] Jacquemart D, Laraia A, Tchana F K, Gamache R, Perrin A and Lacome N 2010 *J. Quant. Spectrosc. Radiat. Transf.* **111** 1209–1222
[109] Cazzoli G and Puzzarini C 2012 *Journal of Quantitative Spectroscopy and Radiative Transfer* **113** 1051–1057
[110] Tasinato N, Charmet A P, Stoppa P, Giorgianni S and Buffa G 2014 *Spectrochimica Acta Part A - Molecular and Biomolecular Spectroscopy* **118** 373–379
[111] Yang C, Buldyreva J, Gordon I E, Rohart F, Cuisset A, Mouret G, Bocquet R and Hindle F 2008 *J. Quant. Spectrosc. Radiat. Transf.* **109** 2857–2868
[112] Devi V M, Benner D C, Smith M A H, Rinsland C P, Sharpe S and Sams R L 2004 *J. Quant. Spectrosc. Radiat. Transf.* **87** 339–366
[113] Rinsland C P, Devi V M, Benner D C, Smith M A H, Sharpe S and Sams R L 2004 *J. Quant. Spectrosc. Radiat. Transf.* **82** 343–362
[114] Devi V M, Benner D C, Smith M A H, Rinsland C P, Sharpe S and Sams R L 2003 *J. Quant. Spectrosc. Radiat. Transf.* **82** 319–342
[115] Mehrotra S, Mader H, de Vreede J and Dijkerman H 1985 *Chemical Physics* **93** 115 – 125 ISSN 0301-0104 URL http://www.sciencedirect.com/science/article/pii/0301010485850539
[116] Hurtmans D, Henry A, Valentin A and Boulet C 2009 *Journal of Molecular Spectroscopy* **254** 126–136
[117] Pine A and Fried A 1985 *Journal of Molecular Spectroscopy* **114** 148–162
[118] Toth R A, Hunt R H and Plyler E K 1970 *The Journal of Chemical Physics* **53** 4304–4307
[119] Rank D H, Eastman D P, Rao B S and Wiggins T A 1963 *Journal of Molecular Spectroscopy* **10** 34–50
[120] Blanquet G, Walrand J and Bouanich J P 1999 *J. Mol. Spectrosc.* **198** 408–415
[121] Misago F, Lepére M and Bouanich J P 2009 *J. Mol. Spectrosc.* **254** 16–19
[122] May R and Webster C 1989 *Journal of Molecular Spectroscopy* **138** 383–397
[123] Waldmann I P, Rocchetto M, Tinetti G, Barton E J, Yurchenko S N and Tennyson J 2015 *Astrophys. J.* **813** 13
[124] Irwin P G J, Teanby N A, de Kok R, Fletcher L N, Howett C J A, Tsang C C C, Wilson C F, Calcutt S B, Nixon C A and Parrish P D 2008 *J. Quant. Spectrosc. Radiat. Transf.* **109** 1136–1150
[125] Barton E J, Hill C, Czurylo M, Li H Y, Hyslop A, Yurchenko S N and Tennyson J 2016 *J. Quant. Spectrosc. Radiat. Transf.*